

ARTHRITIS AUSTRALIA

Arthritis Australia is a not-for-profit organisation that provides support and information for all Australians affected by arthritis.

Contact your State/Territory Arthritis Office to find out about the range of awareness and education programs, support services and resources available.



10 STEPS FOR LIVING WELL WITH ARTHRITIS



ARTHRITIS ACT

Building 18, 170 Haydon Drive
Bruce ACT 2617
PO Box 908 Belconnen ACT 2616

ARTHRITIS NEW SOUTH WALES

Suite 1, 15/32 Delhi Road
North Ryde NSW 2113
Locked Bag 2216 North Ryde NSW 1670

ARTHRITIS NORTHERN TERRITORY

Shop 18, Rapid Creek Business Village
48 Trower Road, Millner NT 0810
PO Box 452 Nightcliff NT 0814

ARTHRITIS QUEENSLAND

Level 1, 42 Costin Street
Fortitude Valley, QLD 4006
PO Box 2121 Windsor QLD 4030

ARTHRITIS SOUTH AUSTRALIA

118-124 Richmond Road
Marleston SA 5033

ARTHRITIS TASMANIA

19A Main Road, Moonah TAS 7009
PO Box 780 Moonah TAS 7009

ARTHRITIS WESTERN AUSTRALIA

17 Lemnos Street
Shenton Park WA 6008
PO Box 34 Wembley WA 6913

ARTHRITIS AUSTRALIA

Level 2, 255 Broadway
Glebe NSW 2037
PO Box 550 Broadway NSW 2007
Phone: 02 9518 4441
Fax: 02 9518 4011
Email: info@arthritisaustralia.com.au

ISBN: 978-0-9805024-3-5-2



Arthritis Infoline: 1800 011 041 www.arthritisaustralia.com.au



MEDICAL AND CONSUMER CONSULTANTS

Lyn Crossley

Arthritis SA
Health Educator

Tanya deKroo

National Resources Coordinator

Mrs Wendy Favorito

Arthritis Australia Consumer Director

Dr Tiffany Gill

President Arthritis SA,
Physiotherapist and Researcher
and Board member

Maggi Koerbin

Arthritis Tasmania

Jacqui Louth

Arthritis NSW

Judith Lowe

Arthritis SA
Health Educator

Dr Mona Marabani

Rheumatologist and President of
Arthritis Australia (2007-2010)

Professor Susanna Proudman

Arthritis Australia Medical Director
and Board member

Jo Rayner

Arthritis SA, Manager Education
Research and Advocacy

Monte Scharbow

Arthritis ACT

Louise Schnitzhofer

Arthritis NT

Monica Tardinai

Consumer

Assoc Professor John York

Department of Rheumatology,
Royal Prince Alfred Hospital Sydney



Proudly supported by an educational grant
from:



© Copyright Arthritis Australia 2019.
Reviewed July 2019

CONTENTS

INTRODUCTION	4
STEP 1 Take control by knowing your disease	5
Arthritis – what is it? • Osteoarthritis • Rheumatoid arthritis • Gout	
STEP 2 Don't delay, see your doctor	9
What to expect • Different types of tests	
STEP 3 Work with your healthcare team	11
Making the most of your healthcare team – some tips	
STEP 4 Know about your treatment options	12
Physical therapies • Medicines – an overview • Complementary therapies	
STEP 5 Find new ways to stay active	17
Why exercise? • What is the right exercise for arthritis?	
STEP 6 Learn techniques to help manage your pain	21
Some pain management techniques	
STEP 7 Acknowledge your feelings and seek support	23
Dealing with depression and understanding your feelings	
STEP 8 Make food choices that count	26
Omega 3 oils • Diet and gout	
STEP 9 Balance your life	29
How to pace yourself	
STEP 10 Call your local State/Territory Arthritis Office	31
Extra info about making informed decisions	32

Introduction

When you have arthritis and are experiencing pain, stiffness or swelling of your joints, everyday tasks around the home may be more difficult.

Arthritis is a very common condition in Australia affecting people of all ages and from all walks of life. Its symptoms often have a big impact on the daily lives of people.

Although arthritis can be difficult to live with, there are many simple measures that can help anyone with arthritis manage their symptoms and cope with daily life. These measures rely on input from a number of people but the most important person is you – the person with arthritis.

This book will put you on the path to managing your arthritis by taking an active role in understanding and treating your condition, both of which will help you to be more active in your daily life.

The key steps listed on this page form a simple-to-follow checklist for living well with arthritis. Read the rest of the information in this book to find out more about each of the steps – and good luck on your journey to better living!

10 STEPS FOR LIVING WELL WITH ARTHRITIS

- 1 Take control by knowing your disease
- 2 Don't delay, see your doctor
- 3 Work with your healthcare team
- 4 Know about your treatment options
- 5 Find ways to stay active
- 6 Learn techniques to help manage your pain
- 7 Acknowledge your feelings and seek support
- 8 Make food choices that count
- 9 Balance your life
- 10 Call your local State/Territory Arthritis Office

STEP 1

Take control by knowing your disease

Learning about your disease is vital.

Spend the time to understand what type of arthritis you have and then discover the best ways to improve your condition.

Many people with arthritis say that learning about their disease helps them to feel in control over their lives and health.

The information in this booklet is a good start to understanding your arthritis.

If you would like more information phone your local State/Territory Arthritis Office or visit www.arthritisaustralia.com.au

See page 31 for contact details of your State/Territory Arthritis Office.

Arthritis – what is it?

Arthritis is not a single disease. The word 'arthritis' is a name for a group of more than 100 diseases that affect the joints. Arthritis can affect many different parts of the joint and nearly every joint in the body. People can be affected in all sorts of different ways but the most common symptoms of arthritis are pain, stiffness and swelling in one or more joints. The three most common types of arthritis are osteoarthritis, rheumatoid arthritis and gout.

Osteoarthritis

The most common form of arthritis is osteoarthritis (OA). It mainly affects people over the age of 40 but it can develop at any age. OA can affect any joint in the body but it usually occurs in large joints that support the weight of your body such as the hips, knees and lower back. It can also occur in the hands, particularly at the base of the thumb and the end joints of the fingers and in the big toe.

It is a condition that affects the whole joint including bone, cartilage, ligaments and muscles.

Osteoarthritis may include:

- inflammation of the tissue around a joint
- damage to joint cartilage – this is the protective cushion on the ends of your bones which allows a joint to move smoothly
- bony spurs growing around the edge of a joint
- deterioration of ligaments (the tough bands that hold your joint together) and tendons (cords that attach muscles to bones).

Osteoarthritis can be a result of:

- an injury to the joint, such as a broken bone or torn cartilage – which might have happened years before arthritis appears

Step 1 Take control by knowing your disease

- being overweight – this puts extra strain on weight-bearing joints and they may become worn under the pressure
- jobs involving repetitive movements, such as heavy lifting, kneeling and squatting
- family history – if your parents have or had arthritis, especially in the hands, you are more likely to develop it.

The symptoms of osteoarthritis vary from person to person. The most common signs are:

- joint pain and tenderness
- stiffness of the joints
- symptoms that worsen after overactivity or underactivity. For example, many people find their symptoms worsen after doing lots of activity such as walking or gardening, or after periods of inactivity such as when getting out of bed in the morning or after sitting for long periods.

Other symptoms may include:

- swelling of the joints
- muscle weakness, which may feel like the joint is unstable or will give way
- a grinding sensation or clicking noises in the joints
- a feeling that the joint might lock
- difficulty walking, using stairs, using your hands.

At present there is no cure for OA, but there are things you can do to manage the pain and limit further damage to the cartilage.

One of the most important things you can do is exercise the joints. The joint cartilage relies on the fluid in the joint ('synovial fluid') to keep it healthy by providing nutrients and lubrication. Movement encourages the body to produce synovial fluid and keeps it moving around the entire joint. Exercise also strengthens the muscles that support the joints and gives you greater mobility and flexibility.

[See page 17 for more details about exercise.](#)

Arthritis Australia has developed a website www.MyJointPain.org.au to help people with osteoarthritis. The site can help create a tailored OA management plan for you. It provides advice about OA and tools to help you manage symptoms. It can also help you find support from local healthcare professionals.

Rheumatoid arthritis

Rheumatoid arthritis (RA) is a disease that causes inflammation of the joints. Any joint can be affected however it often affects smaller joints such as the hands and feet. Larger joints such as the hips and knees can also be affected.

Women are three times more likely to develop RA than men and it can happen at any age. Rheumatoid

arthritis is classified as an autoimmune disease. The normal role of your body's immune system is to fight off infections to keep you healthy. In an autoimmune disease, your immune system starts attacking your own healthy tissues. In RA, the immune system targets the lining of the joints ('synovial membrane'), which becomes inflamed and thickened. This causes the body to produce larger than normal amounts of fluid in the joints ('synovial fluid'), which leads to swelling, pain and stiffness.


If the joint continues to be inflamed, the bones, the cartilage that covers the ends of the bones, and the muscles and ligaments of the joint can all become damaged. This inflammation and swelling can cause the joint to become unstable, making it difficult to use the joint.

The symptoms of rheumatoid arthritis vary from person to person but often include:

- joint pain and tenderness
- early morning stiffness in the joints
- swelling in the joints
- the same joints on both sides of the body are affected (ie symptoms in both hands or both knees).

Currently there is no cure for rheumatoid arthritis but there is a lot you can do to control it, manage the pain and live a full life with the disease:

- see your doctor immediately – early diagnosis and treatment



can limit joint damage caused by inflammation

- learn pain management techniques – like relaxation, mindfulness or using hot and cold packs
- be physically active – consider warm water exercise, strength training and tai chi – these will strengthen supporting muscles and keep your joints flexible. They will also maintain your feeling of wellbeing
- learn how to protect your joints and pace yourself – use equipment such as specially designed cooking utensils and take breaks when needed. This will help you reduce stress on the joints and help you manage fatigue
- stop smoking – smokers are twice as likely to get RA as non-smokers and smoking may effect some of the medications prescribed for RA
- take the medicines recommended by your doctor
- seek support from other people and support groups – for advice, social contact and emotional support.

Step 1 Take control by knowing your disease

Arthritis Australia has developed a website that brings together a range of people living with rheumatoid arthritis and other similar conditions. To hear directly from them, and their medical team, on how they have managed to survive and thrive visit www.empowered.org.au

Gout

Gout is a common and painful condition. There are many myths about the causes of gout. Gout is not caused by over eating and drinking too much alcohol but occurs as a result of high levels of uric acid in the body. In fact, gout can affect anyone.

Uric acid is a normal waste product but people with gout have higher-than-normal amounts of uric acid in their bodies. If uric acid levels remain high, tiny crystals form in the joints and other organs of the body. This causes inflammation, pain and swelling.

Unlike other types of arthritis, which develop slowly, an attack of gout can occur quickly, even overnight. While gout often affects the joint of the big toe it can occur in any joint. Knees, ankles and hands are other common joints affected.

Even if an attack has finished, gout does not go away. If the causes of gout are left untreated, attacks may become more common and the joints can become permanently damaged.

See your doctor to help manage an attack and also prevent future attacks.

Gout is aggravated by anything that reduces the kidneys' ability to remove uric acid from the blood.

Gout attacks can be prevented by:

- taking medicines to lower uric acid levels in your blood. These medicines need to be taken every day, whether you are having an attack or not
- cutting down the amount of alcohol you drink and avoid drinking a lot of alcohol at one time (binge drinking). Talk to your doctor or see www.alcohol.gov.au for Australian Government guidelines on recommended alcohol intake
- keeping a healthy body weight. An obese person is four times more likely to develop gout than someone with ideal body weight. However, it is important to avoid fasting or 'crash' dieting as this can actually increase uric acid levels and trigger a gout attack. A combination of healthy eating and regular physical activity is the best approach
- having a healthy balanced diet.

See page 28 for more information about diet and gout.

Step 2

Don't delay, see your doctor

Early diagnosis of arthritis provides the best treatment outcomes.

This will help you understand your arthritis and develop a plan for managing it. Early diagnosis and treatment can limit the effects of arthritis on your life and help you stay active and independent.

What to expect when you go to the doctor

When you first visit your doctor, you will be asked a number of questions about your symptoms including:

- how long you have experienced pain
- which joints are affected
- when you get pain and what seems to cause it
- what makes the joint feel better or worse
- whether anyone else in your family has had arthritis or joint pain.

Your doctor may also order tests including:

- blood tests
- a test of your joint fluid ('synovial fluid')
- x-rays.


At your next appointment, your doctor will discuss the results with you but it may take several visits to your doctor before the type of arthritis is diagnosed. This is because there is often no single test to diagnose individual types of arthritis and the symptoms can be similar.

What are the different types of tests

Blood tests and pathology tests

There are several blood tests your doctor may order. The main ones used to check for common types of arthritis are:

- uric acid levels – a higher than normal level of uric acid means you may have gout
- antibody tests – antibodies are substances the body normally produces to fight infections. In some forms of arthritis, the body produces specific antibodies, for example seven out of ten people with rheumatoid arthritis have a 'rheumatoid factor'



- inflammation tests – there are two main tests used to check if you have inflammation: ESR (erythrocyte sedimentation rate) and CRP (C-reactive protein). Together with the other blood tests, your ESR and CRP levels will help your doctor decide whether you have a form of inflammatory arthritis such as rheumatoid arthritis.

Sometimes your doctor may order other, more specialised blood or pathology tests.

At the same time as you have tests for arthritis, your doctor may also order other tests to check your general health.

Tests of the joint fluid (synovial fluid)

To help diagnose some types of arthritis, your doctor may take a sample of fluid from the affected joint.

Your doctor will do this by putting a needle directly into the joint, then drawing out a small amount of the fluid. It can hurt slightly but local anaesthetic may be used before the procedure. Your doctor will then have the sample checked under a microscope to see if crystals, such as uric acid from gout, are present.

X-rays

X-rays provide pictures of bones and can show changes in the shape of the joint, such as narrowing or deformities. X-rays are not always helpful in the early stages of arthritis as changes in the joint structure may not be visible for months or even years after arthritis begins. The amount of pain you are experiencing may not equal the amount of damage shown on the x-ray. For example, your joint may be very painful despite x-rays being normal. Pain can be associated with soft tissues such as muscles or tendons that are not visible on an x-ray. Your doctor will only order an x-ray if the cause of your symptoms is not clear, if your symptoms are severe or have become a lot worse.

Step 3 Work with your healthcare team

The best way to live well with arthritis is by working closely with your healthcare team.

It may include a variety of healthcare specialists, such as doctors, pharmacists, physiotherapists, occupational therapists, podiatrists, exercise physiologists, nurses, psychologists and complementary medicine practitioners.

Your doctor may also refer you to a rheumatologist (a specialist in arthritis) who can provide you with the most up-to-date information and advice on all aspects of arthritis management and treatment.

Remember, you are the most important member of your healthcare team. Make sure you establish and maintain good communication with all the other members.

Making the most of your healthcare team

To get the most out of a visit to your doctor or other healthcare professional it is important to be prepared. Here are some tips:

- Think about, and write down, the questions you want to ask before your visit

- Always take test results related to your condition to the consultation
- Consider taking a family member or friend with you as a second set of ears
- Ask your doctor or healthcare professional to explain any information that you did not understand
- Feel free to ask questions, especially about the benefits, side effects and costs of treatments
- Tell your doctor or healthcare professional if you need time to think or to discuss something with family members
- To help you remember any important information or instructions you are given, write it down or ask for a leaflet or website to read later
- Ask your doctor or healthcare professional where you can learn more about your condition or treatment
- Tell your doctor about any other medicines, herbal remedies, creams or any other products you are using - including any you have bought from a supermarket, pharmacy or health food store. Any of these may interact with your arthritis medicines causing serious health problems or make other medicines (e.g. prescription medicines) less effective.

If you are unsure about treating your arthritis or planning for the future, discuss this with one of your healthcare team or seek a second opinion. They can also advise you about support services for people with arthritis and where to find more information.

Step 4

Know about your treatment options

There are many treatments to relieve pain and stiffness and slow the development of your arthritis.

Work with your healthcare team to find a combination of treatments that best suits:

- your type of arthritis
- the joints affected
- the amount of pain or other symptoms you experience
- your lifestyle.

A treatment program may include a combination of:

- exercise
- weight loss and/or healthy eating
- physical therapies including physiotherapy, occupational therapy and podiatry
- pain management techniques such as relaxation and meditation
- medicines including prescription, non-prescription and complementary medicines
- emotional and social support
- finding a balance between activity and rest.

See page 22 of this book for professional associations

Physical therapies

Physical therapies are often an important part of a successful arthritis treatment program.

Physiotherapists, occupational therapists, podiatrists and other therapists offer a range of treatments that will help your joints function better.

When choosing any kind of therapist, look for an accredited practitioner or one who is a member of their professional association.

Physiotherapy

Physiotherapists can advise you on exercise, posture and non-medicine-based pain relief. They may also use techniques to keep your joints and muscles flexible.

Treatments that physiotherapists may use include:

- joint mobilisation
- hydrotherapy – exercise in water
- electrotherapy – the use of mild electrical impulses to treat pain and swelling
- braces and taping
- muscle strengthening exercises and stretches.

Occupational therapy

Occupational therapists can advise you on how to take stress and strain off joints affected by arthritis.

They look at all aspects of your daily life, including your job, the work you do around your home, as well as your leisure activities. They can then show you ways to conserve your energy by simplifying daily tasks and how to protect your joints when you are performing those tasks.

Occupational therapists can also advise you on:

- any special equipment you might need to help you, such as grab rails, walking sticks or wheelchairs
- splints and braces you might need, particularly for your hands
- aids and equipment that can make daily activities such as showering, cooking, and driving easier on your joints.

Podiatry

Podiatrists treat problems in the feet. They can help you with footwear, nail care and orthoses (shoe inserts) if you have arthritis in your feet, legs or lower spine.

Medicines

Medicines can help:

- relieve arthritis symptoms, such as pain
- reduce inflammation and swelling
- slow down damage to joints in some types of arthritis.

The medicines your doctor or rheumatologist prescribes will depend on your type of arthritis and the severity of your symptoms.

The most common examples include:

- pain relievers
- non-steroidal anti-inflammatory drugs
- corticosteroids and injections into the joint
- disease-modifying medicines.

Always talk to your doctor and pharmacist before you start taking any medicines.

Find out what side effects may occur and make sure you know what to do if you develop any side effects and how quickly you need to act.

See page 32 for more information on how to take arthritis medicines safely.

Pain relievers (analgesics)

Pain relieving medicines have no effect on the joint inflammation or your arthritis, they simply stop you feeling the pain. Paracetamol is often used for the relief of all types of musculoskeletal pain including arthritis. Unlike ibuprofen (brand name Nurofen) and other non-steroidal anti-inflammatories (NSAIDs), paracetamol is not effective in reducing the inflammation associated with arthritis. You may wish to try paracetamol for a short period to see if it works for your pain. Ask your pharmacist or doctor for advice.

Step 4 Know about your treatment options

For severe, uncontrolled pain your doctor may prescribe stronger pain relievers.

Non-steroidal anti-inflammatory drugs

Non-steroidal anti-inflammatory drugs are commonly referred to as 'NSAIDs' (pronounced 'ensayeds'). NSAIDs reduce inflammation, joint swelling and stiffness. They are often used to treat inflammatory forms of arthritis, such as rheumatoid arthritis. They can also relieve pain that is not controlled by analgesics alone. Some NSAIDs are available without a prescription, while others must be prescribed by a doctor.

NSAIDs stop the body producing substances that cause inflammation and pain. There are many different types of NSAIDs available, such as naproxen, ibuprofen or aspirin. There is also a certain group of NSAIDs called COX-2 inhibitors that are slightly less likely to cause stomach problems.

There has been a lot of controversy recently about anti-inflammatory drugs. Some studies have suggested that there may be a small increase in risk of heart attack or stroke in people who take these drugs at high doses. People with high blood pressure, kidney or heart problems or at risk of stomach bleeds need careful assessment by their doctor to make sure they are on the right drug at the most suitable dose

for the appropriate length of time. Most people will have no problem but you should talk to your doctor or pharmacist before taking these medicines.

Corticosteroids

If you have severe pain and inflammation in your joints, then your doctor may prescribe a stronger anti-inflammatory medicine called a corticosteroid. These can be taken as tablets or given by injection directly into a joint, muscle or other soft tissue.

Corticosteroids can have serious side effects if taken in high doses or for a long time (more than a few weeks). Your doctor will closely monitor you for side effects while you are taking corticosteroids.

Disease-modifying medicines

These medicines are used for inflammatory forms of arthritis, such as rheumatoid arthritis. Discuss these highly specialised medicines with your rheumatologist.

Complementary therapies

'Complementary therapies' are any treatments or therapies that are not part of the conventional treatment (such as medicines or surgery) of a disease. Complementary therapies are taken alongside conventional treatments and may also be known as 'traditional' or 'alternative' therapies. Examples of complementary therapies



include acupuncture, massage, aromatherapy, naturopathy and herbal medicines.

Just like using conventional treatments, you should take reasonable care when using complementary medicines and therapies, including getting your doctor's advice. Not all complementary therapies are regulated and may not be standardised in terms of purity, dosage, effectiveness or safety.

This does not mean you should not try complementary therapies but make sure you are fully informed about the treatment and seek advice from a qualified practitioner so that you are not misled or given false hope.

Like any other treatment, complementary therapies may not be suitable for everyone – nutritional supplements may interact with prescribed medicines causing undesirable effects; magnetic

therapy may cause interference with pacemakers; and some herbs and supplements should not be taken during pregnancy or if you are intending to get pregnant.

For information about registered complementary therapy practitioners, contact the Australian Traditional Medicine Society on 1800 456 855, visit www.atms.com.au or contact the relevant registration body.

Weight control

Being overweight is a risk factor for some forms of arthritis, especially osteoarthritis (OA). Weight loss has been shown to significantly lessen pain in some joints and to help mobility and joint function. A dietitian and exercise physiologist can help devise a weight management plan and assist you with this.

See page 26 for more information

Step 4 Know about your treatment options

Joint replacement

If other treatments have not worked for you, and you are unable to go about your day-to-day life because of pain and loss of mobility, you may require joint-replacement surgery. These are commonly performed operations and most people find that joint replacement surgery improves their quality of life.

Your doctor will refer you to an orthopaedic surgeon who can advise you on whether surgery is required.

To make a fully informed choice about surgery, find out about the possible risks of surgery, waiting lists, costs and recovery time.

Glucosamine and chondroitin

Glucosamine and chondroitin are substances naturally produced by the body and may be useful for people with osteoarthritis, where there has been a breakdown of cartilage. It is thought that taking glucosamine and/or chondroitin supplements may relieve the pain and prevent or slow the breakdown of cartilage in OA.

Note, most of the studies have looked only at OA of the knee, with very few studies of other joints (for example, hips, hands, back). To date there is no evidence that these supplements are effective for any other forms of arthritis.

See Arthritis Australia's [glucosamine and chondroitin information sheet for more about these supplements.](#)

As with all supplements, you should ask your doctor or pharmacist for advice and information about possible side effects before taking glucosamine and chondroitin and check whether you can take it with your other medicines.



Step 5 Find new ways to stay active

Research has found that regular exercise is one of the most effective treatments for arthritis.

Physical activity helps in two ways.

Firstly, exercise will help your arthritis by:

- decreasing the pain in your joints and muscles
- maintaining and increasing the flexibility of your joints and muscles
- strengthening muscles – this will help take the load off your joints, and make the joints more stable
- decreasing or relieving muscle tension – tension adds to the pain of arthritis and in the long term can lead to poor posture and joint deformity
- improving your posture and balance to reduce your risk of falling

Secondly, regular exercise will improve your overall health.

This means you will be able to do more in life, feel more in control of your arthritis and be better able to manage pain.

In particular, exercise will:

- improve your heart and lung fitness
- help control weight and reduce body fat – this will also reduce the load on weight-bearing joints such as feet, knees and hips
- decrease stress and depression
- strengthen bones
- improve your sleep
- decrease fatigue and tiredness
- create a feeling of general wellbeing.

Types of exercise that are beneficial for arthritis

Not all forms of exercise are appropriate for every kind of arthritis. Before you start to exercise, it is important to ask your doctor and healthcare team to help you develop a program that will suit your type of arthritis, general health and lifestyle.

Generally, you will need to do a mix of three types of activities:

- mobility exercises
- strengthening exercises
- fitness exercises.

Step 5 Find new ways to stay active

Mobility

These are exercises designed to maintain or improve the range of movement of the joints and muscles.

Mobility exercises are especially important for stiff joints and muscles, although all your joints will benefit from maintaining or improving their range of movement.

Examples include moving each joint as far as it will comfortably go, through muscle stretches or yoga.

Strengthening

These exercises are designed to increase the power of the muscles. Strong muscles help to support and take pressure off sore joints. Strengthening exercises also help strengthen bones and improve balance.

Examples include exercises using weights, dumbbells and resistance bands.

Fitness

These are the exercises that will benefit the heart, lungs and your general wellbeing. These are usually activities that use the larger muscles in the body, rather than exercising a specific area, and may make you 'puff' a little. Examples include brisk walking, swimming and cycling.

Arthritis-friendly physical activity

There are many activities that can be beneficial for people with arthritis.

The best activities are those you find enjoyable and are convenient. Low-impact exercises, with less body weight or force going through your joints, are usually most comfortable.

Examples of low-impact activities include:

- walking – a simple way to increase fitness
- exercising in water, such as hydrotherapy (with a physiotherapist), swimming or water exercise classes
- strength training – using weights, resistance bands, gym machines or even just your own body weight
- tai chi, yoga and pilates – good for flexibility and strength, as well as relaxation and stress management
- cycling
- dancing – excellent for flexibility and fitness and also helps build stability in the joints
- chair-based exercises.

Water exercise is a popular exercise for people with arthritis. The buoyancy of the water takes pressure off painful joints and you may find you can move more freely than you can on land. Warm water can also be soothing for sore muscles and stiff joints.



Getting started

Talk to your doctor and/or health professional before starting an exercise program. A physiotherapist or exercise physiologist can suggest safe exercises and make sure you are doing your exercises correctly to prevent an injury.

As a general guide, all Australian adults should be aiming to do at least 30 minutes of activity on most days of the week. You can do 30 minutes continuously or combine several 10 to 15 minute sessions. If you have arthritis and you have not exercised for a while, you may need to start with shorter sessions (for example, 5 to 10 minutes). When you first start, do less than you think you will be able to manage. If you cope well, do a little bit more next time and keep building gradually.

Always start your exercise with some gentle movements to warm up your body and your joints. This can help prevent pain and injury during exercise.

Cool down at the end of your session with some gentle movements and stretches. This can help prevent muscle pain and stiffness the next day. Your local Arthritis office can also connect you to local arthritis-friendly physical activity programs.

If the thought of 'exercise' seems too hard, try increasing your daily physical activity. For example:

- get off the bus one or two stops earlier than you normally would and walk the rest of the way
- walk up stairs instead of waiting for the lift, or walk up one flight of stairs then take the lift
- park the car a little bit further from the shops instead of looking for the closest parking space.

Sticking to an exercise program

If you have not been active for a while or you have pain and often feel tired and unwell, it can be difficult to make exercise a regular part of your life.

Here are some tips for sticking with an exercise program:

- do activities you enjoy
- do something that is convenient and affordable
- exercise with a friend or in a group
- start gradually and build up
- set short-term goals for what you want to achieve. Write them down and put them on the fridge
- keep an exercise diary, so you can see how you are progressing
- vary your exercise routine so you don't get bored.

When to exercise

Try to do some form of exercise every day, even on bad days. If possible, try to exercise when:

- you have least pain
- you are least stiff
- you are least tired, and
- your medicines are having the most effect (ask your doctor or pharmacist about how to time your medicines with exercise if possible. This may help to make your exercise session more comfortable).

When to stop exercising or be careful

- Don't vigorously exercise a joint that is red, hot, swollen or painful
- Stop exercising if it is causing you unusual pain or increases your pain beyond what is normal for you.

Exercising through this type of pain may lead to injury or worsening of your arthritis symptoms. (Note, many people with arthritis have some amount of pain all the time. This is not a reason to avoid exercise. You should only stop if you notice extra or unusual pain while you are exercising)

- Use the two hour pain rule to measure how well your body is coping with an activity – if you have extra or unusual pain for more than two hours after exercising, you've done too much. Next time you exercise, slow down or do less
- Talk to your doctor, exercise physiologist or physiotherapist about any problems or questions you have about your exercise routine.



STEP 6 Learn techniques to help manage your pain

You may have to accept that sometimes medicines, physical therapies and other treatments cannot relieve all of your pain.

Pain may limit some of the things you do, but it doesn't have to control your life. There are many techniques you can use to cope with pain so you can go on living your life the way you want to. Your mind plays an important role in how you feel pain. Thinking of pain as a signal to take positive action rather than being scared or worried about it can be helpful. Also you can learn ways to manage your pain. What works for one person may not work for another, so you may have to try different techniques until you find what works best for you.

Here are some things you can do to manage your arthritis pain:

- make the most of your medicines and physical therapies. Ask questions of your healthcare team and discuss any concerns. Visit your doctor regularly to make sure you are getting the best treatment for your arthritis symptoms
- take care of your body. Exercise to improve your fitness and strength, eat a healthy diet, aim to get a good night's sleep every night

- use heat and cold treatments for extra pain relief. A warm bath or shower, or a heat pack placed over a painful joint for 15 minutes, can provide effective pain relief. An ice pack may reduce swelling and relieve pain in the same way. Ask your doctor or physiotherapist which type of treatment (hot or cold) is best for you
- find some distraction techniques that work for you. These may include exercising, reading, listening to music, or seeing a movie. Anything that focuses your attention on something enjoyable, instead of your pain, will help you
- learn some relaxation techniques. When you are stressed, your muscles become tense, making pain feel more severe. Relaxation techniques such as meditation or deep breathing help to decrease muscle tension
- ask your physiotherapist about transcutaneous electrical nerve stimulation (TENS). A TENS machine applies very mild electric pulses via small electrodes (pads) to block pain messages going from the painful area to your brain. TENS



STEP 7 Acknowledge your feelings and seek support

It is natural to feel frustrated, angry, scared, anxious and even depressed at the prospect of having arthritis.

Many people with arthritis fear what the future might hold and are frightened by the impact arthritis might have on their everyday life. It is not unusual either for a young person to feel especially angry or depressed at being diagnosed with a disease that is mistakenly thought to affect only 'old' people.

How to cope

Arthritis can occur at any age including in children and young adults. It is not great news to find out you have a chronic health condition and it is not surprising that people can have negative thoughts and emotions such as a sense of loss or distress.

To help you cope with the changes that arthritis brings you need to:

- find new activities and challenges that you can do with arthritis and that will give you satisfaction
- think about this process of adjustment as "just another part of growing" – it is about learning new skills and adjusting to experiences

- talk to people who have gone through what you are experiencing. Their experiences may not be identical but they can tell you about things that helped them
- talk to your friends, your doctor, a counsellor or a psychologist if you have strong feelings of loss or unhappiness. You are not alone and do not have to make these adjustments all by yourself. There are always people who can listen to you and help. Most people find talking through their experiences helpful.

Stress

Stress is the mind/body's response to perceived fear or pressure. When you are in a demanding situation, you may have worrying thoughts and your brain releases chemicals that stimulate the rest of your body; your heart beats faster, your breathing changes, blood pressure rises and muscles become tense. People with arthritis need to learn how to manage stress, otherwise it can lead to muscle tension, pain and depression.

can be useful for relieving pain but does not work for all people. You should see a physiotherapist to trial a TENS machine, and to learn how to use it correctly

- some people with arthritis find massage and acupuncture useful to help control pain and improve relaxation. It is wise to seek a registered practitioner who understands your condition.

You can find qualified massage therapists by contacting **Massage and Myotherapy Australia (formerly the Australian Association of Massage Therapists)** at www.aamt.com.au or (03) 9602 7300 or the **Myotherapy Association of Australia** at www.massagemyotherapy.com.au or (03) 9418 3913.

The **Australian Acupuncture and Chinese Medicine Association** can help you find an **accredited acupuncturist** at www.acupuncture.org.au or 1300 725 334.

Step 7 Acknowledge your feelings and seek support

Managing stress involves two steps:

1. recognising when you are becoming stressed – learn to listen to your body and recognise the signs
2. finding activities that will relieve stress.

There are many ways you can reduce stress. Talk to your doctor and other members of your healthcare team to find solutions that work for you.

Some things that you can do for yourself that will help manage stress include:

- being physically and mentally active – but don't overdo it to the point of fatigue
- learning relaxation techniques – like meditation, yoga and tai chi
- learning deep breathing techniques – slowing and deepening your breathing will counteract the body's stress response of fast, shallow breathing. Slow, deep breathing can also slow your heart rate
- talking to friends and your healthcare team about things you find stressful – researchers have found that some people also benefit from writing about stressful experiences
- learning what situations you find stressful and either avoiding them, changing them or learning how to use your relaxation techniques when they occur

- getting support from friends, family or other people with arthritis.

Dealing with depression and anxiety

Depression is not just a low mood or feeling sad, but a serious condition that needs treatment. People with depression generally feel sad, down or miserable most of the time. They find it hard to be interested in normal day-to-day activities. Depression has serious effects on physical as well as mental health.

Depression is common and affects up to one million people in Australia each year. Signs to watch for are:

- feeling uncontrollably sad
- withdrawing from friends and family
- losing interest and pleasure in most of your usual activities
- thinking that no-one understands what you are going through
- feeling unable to cope, even with everyday things
- losing or gaining weight
- disturbed sleeping patterns
- tiredness and lack of energy
- poor concentration and not thinking clearly.

Like any other illness, the sooner you seek help the quicker your depression can be treated. Do not be afraid or embarrassed to speak with your doctor or healthcare team.

They will be able to reassure you that your feelings are not unusual and that depression can be treated safely and effectively.

Treatment for depression will depend on how long you have been depressed and the severity of your depression and may include:

- medicines
- counselling
- consultation with a psychiatrist.

Anxiety is more than just feeling stressed or worried. Anxiety is when these anxious feelings don't settle down – when they are ongoing and are there for no particular reason or cause. It's a serious condition that makes it hard to cope with daily life. Everyone feels anxious from time to time but, for someone with anxiety, these feelings can't be easily controlled.

Many people with anxiety may experience depression as well. Having arthritis is one of many factors that may trigger anxiety. Anxiety is common and the sooner you get help, the sooner you can begin to recover.

While everyone's experience with anxiety will be different, there are some common symptoms including:

- Physical: panic attacks, hot and cold flushes, racing heart, tightening of the chest, quick breathing, restlessness, or feeling tense, wound up and edgy
- Psychological: excessive fear, worry, catastrophising, or obsessive thinking

- Behavioural: avoidance of situations that make you feel anxious which can impact on study, work or social life.

If you are experiencing symptoms of depression or anxiety, there is help available. Find someone you feel comfortable talking to and express your feelings. This could be a member of your family, a friend, your doctor, a psychologist, counsellor or a support group.

Contact your local State/Territory Arthritis Office to find out more about support services for people with arthritis or speak to your doctor.

beyondblue provides information and advice about depression, anxiety, available treatments and where to get help. Visit www.beyondblue.org.au or call 1300 22 4636.

Lifeline provides a 24hr confidential telephone crisis support service for anyone across Australia needing urgent help. Call 13 11 14.

Step 8

Make food choices that count

Food and arthritis

There are many myths about food and arthritis. No diet has been proven by research to cure arthritis and there is very little scientific evidence that specific foods have an effect on arthritis. For example, it is unproven that 'acidic' foods such as tomatoes can cause arthritis or make it worse. Except for gout, most foods will not have a direct effect on your arthritis, or on the pain and stiffness you experience.

- Be very cautious of special diets or supplements that claim to 'cure' arthritis or control its symptoms
- A healthy, balanced diet is important for maintaining your general health and wellbeing
- If you are overweight, a healthy diet will also help you reduce your weight, which will lessen the pressure on weight-bearing joints such as knees, ankles, hips and spine. This will help reduce the amount of pain you experience
- Some people with certain forms of arthritis can lose their appetite. If this happens to you, try eating more meals in smaller portions throughout the day. This will help provide the sustenance you need.

For more information about general healthy eating guidelines see the Australian Government's Australian Dietary Guidelines at www.eatforhealth.gov.au

For more specialised advice, ie if you have gout, on how to have a balanced diet or healthy weight loss, talk to a dietitian. You can ask your GP to refer you to a dietitian or find one directly via the Dietitians Association of Australia call 1800 812 942 or visit www.daa.asn.au

There is also information and practical tips about weight control at www.8700.com.au

Foods that affect and reduce inflammation

Omega 3 oils

Although most foods have no effect on arthritis, studies show that eating foods rich in omega-3 fats can help reduce inflammation in some forms of arthritis. While these effects are modest compared to medicines, omega-3 fats do not have serious side effects.

Current research suggests omega-3 fats are helpful for people with inflammatory arthritis, such as rheumatoid arthritis.

Foods rich in omega-3 fats:

- oily fish: sardines and salmon have greater amounts of omega-3 fats. Try to eat these types of fish at least two to three times a week
- fish oil supplements: see the section below for more information
- ground linseeds and linseed oil (also called flaxseed)
- walnuts
- canola oil (also called rapeseed oil).

Fish oil supplements

Fish oil supplements are available as capsules or as a liquid. Different brands of capsules differ in the amount of omega-3 fats they contain so it is worthwhile to compare brands. Research suggests the dose needed to reduce inflammation in conditions such as rheumatoid arthritis is 2.7 grams of omega-3 (EPA plus DHA) daily.

This dose usually requires either:

- nine to 14 standard 1000mg fish oil capsules or five to seven capsules of a fish oil concentrate per day, or
- 15mL of bottled fish oil or five to seven ml of concentrated bottled fish oil per day.



For osteoarthritis, fish oil supplements at a lower dose (providing 0.45 grams of omega-3) may be useful. You may need to take fish oil supplements regularly at the recommended arthritis dose for two to three months before you notice improvements in your arthritis symptoms. If there is no change by then, the supplements are probably not effective for your arthritis.

It is important not to confuse fish oils with fish liver oils (such as cod liver oil and halibut liver oil). Fish liver oils also contain vitamin A. Large amounts of vitamin A can cause serious side effects, particularly during pregnancy. To increase your intake of omega-3 fats, you should do so by taking pure fish oils, not fish liver oils.

As with all supplements, you should ask your doctor or pharmacist for advice before taking fish oil and check whether you can take it with your medicines.

Diet and gout

Some foods are commonly linked to attacks of gout. Often these foods tend to contain high levels of purines, a substance that can be made into uric acid in the body.

Purine-rich foods include:

- meat – particularly red meat and offal, such as liver, kidneys and heart
- seafood – particularly shellfish, scallops, mussels, herring, mackerel, sardines and anchovies
- foods containing yeast – such as Vegemite and beer.

Avoiding these foods may not prevent an attack of gout and you may miss out on important nutrients and vitamins by completely cutting these foods from your diet.

You should get tailored advice from your doctor or a dietitian before starting a special diet for gout.

For most people with gout, a healthy balanced diet is all that is needed to manage gout, alongside medicines to reduce uric acid levels. If you notice certain foods trigger your gout attacks, you may benefit from cutting down the amounts of those foods in your diet. However not all purine-rich foods are linked to gout.

For example, a number of vegetables (asparagus, mushrooms, cauliflower and spinach) are rich in purines but do not appear to trigger gout attacks.

Another food source that can increase uric acid levels in the blood and lead to gout attacks is a type of sugar called fructose. Fructose is a sugar that is found in fruits and vegetables. It is also found in high levels in foods sweetened with corn syrup, such as bread, cereal, some soft drinks (not Australian-made soft drinks) and fruit juices. However there is no research showing that fructose actually causes gout.

Cutting down the amount of food artificially sweetened with high fructose corn syrup may be beneficial for your overall health. However naturally occurring fructose in fruit and vegetables also provides general health benefits and should not be completely avoided without advice from your doctor or dietitian.

You should get tailored advice from your doctor or a dietitian before starting a special diet for gout.

Step 9 Balance your life

When you have arthritis, you need to find the right balance between work, activity and rest.

Many people find arthritis a tiring disease particularly if they are in pain. If you find this to be the case, listen to your body and be guided by it.

Rest when you are tired and don't force yourself to work or exercise through pain.

Learn how to pace yourself

Carefully plan and organise your activities so you make the most of your energy.

Here are some simple tips to help you pace your activities:

- Try to plan your day so that you can alternate periods of activity with periods of rest
- When you know you have a large task to do, such as preparing a meal or cleaning a room, plan ahead and break the job into smaller tasks. Then work on completing the tasks one at a time, and follow each with a rest break
- Try to prioritise jobs. Do the hardest jobs when you are feeling your best
- Take advantage of 'good days' to do the things you may have been putting off. Remember not to overdo it on these days as it could result in pain and fatigue the following days

- Simplify tasks. For example, buy pre-cut vegetables and meat to make cooking simpler. Find out about appliances that can make tasks easier (see below)
- If you are having a bad day, be ready to change your plans and not force yourself to work through pain
- Ask for help when you need it
- Visit an occupational therapist to discover ways you can save time and energy.

Learn about equipment that can make daily tasks easier

There are many appliances that can make your daily activities simpler and less tiring. These aids and equipment aim to protect your joints by reducing the effort you have to put in.

Examples include:

- adapted cutlery and cooking utensils to allow easy gripping
- tools to help with opening jars or bottles, and turning on taps
- equipment to make dressing and showering easier (such as long-handled sponges for reaching your feet, long-handled shoe horns).



You can find out more information about these types of equipment at an Independent Living Centre. The centres do not sell items but are a one-stop reference for discovering what is available and where you can buy it. You can get advice, including where to purchase equipment, in person or over the phone.

Occupational therapists are also available at the centres to provide advice about equipment. Independent Living Centres are located in each capital city. See www.ilcaustralia.org.au or call 1300 885 886 to find your closest centre and more information. (Note, the Independent Living Centre is called LifeTec in QLD and Assistive Technology Australia in NSW).

Step 10 Call your local state/territory arthritis office

You can have arthritis and still get the most out of life. You don't have to stop doing the things you love doing.

Your local State/Territory Arthritis Office offers a range of services which may include a telephone information service, regular information sessions and arthritis self-management courses, online webinars, and support groups.

These services will introduce you to a wide range of skills and small changes you can make that can lessen the impact of arthritis on your life, including:

- how to manage symptoms such as fatigue and depression
- how to communicate effectively with your doctor and healthcare team
- how to fight fatigue
- how to get more out of life.

Phone your local State/Territory Arthritis Office now on 1800 011 041 to find out more about the services they offer.

CONGRATULATIONS!

Now that you've learned more about living well with arthritis, you are already on your way to successfully managing your arthritis and lessening its impact on your life. It's important to put what you have learned into practice.

Re-visit the arthritis check list on page 4 from time to time to ensure that you continue to maintain control over your arthritis.

Extra info about making informed decisions

Medicines

All medicines have risks and benefits.

Always talk to your doctor or pharmacist about the medicines you are taking, including supplements, any herbal or complementary treatments, and what you are using them for, so they can help you choose the most effective medicine for your condition, health and lifestyle.

Before you take any kind of medicine

Tell your doctor if you have allergies to:

- any medicines – particularly aspirin, painkillers or anti-inflammatory medicines
- any other substances – such as foods, preservatives or dyes.

If you are pregnant, intend to become pregnant or breastfeeding, talk to your doctor as some medicines used to treat pain and inflammation are not recommended during these times. Inform your doctor of any other medical conditions you have because they may affect the medicines you can take and the best dose for you.

Supplements, herbal or complementary medicines

It is important to tell your doctor about all of the other medicines you are taking. This includes any that you buy without a prescription from a pharmacy, supermarket or health food shop. Some treatments may interfere with arthritis medications.

What Can I do?

Read the Consumer Medicine Information leaflet

Consumer Medicine Information (CMI) leaflets are available for all prescription medicines and some medicines that you buy over-the-counter at a pharmacy. Ask your pharmacist or doctor for a Consumer Medicine Information leaflet for your medicine.

CMI leaflets provide easy to understand information on:

- what the medicine is for
- how to use the medicine
- possible side effects and what to do if they occur.

Call the Medicines Line Medicines Line is a free telephone service

Phone: 1300 633 424

Monday to Friday,

9am to 5pm AEST

www.nps.org.au

Read the Australian Rheumatology Association's patient medicine information on the Arthritis Australia website

The Australian Rheumatology Association has developed a series of information sheets in collaboration with Arthritis Australia, covering most of the medicines commonly used to treat arthritis. Copies can be obtained by ringing the Arthritis Infoline 1800 011 041 or visiting www.rheumatology.org.au or www.arthritisaustralia.com.au

TREATMENT

Not all information you read or hear is trustworthy so always talk to your doctor or healthcare team about treatments you are thinking about trying.

- Get an accurate diagnosis from your doctor. It is important to know exactly what type of arthritis you have before you start any treatment
- Get information about the medications and treatments recommended. Talk to your doctor or pharmacist
- Ask about costs. Make sure the treatment or therapy is something you can afford, particularly if you need to keep using it
- Check a practitioner's qualifications. Most therapies have a professional association that you can contact for more information or they can help you find an accredited practitioner.

Some qualifications can be checked at www.ahpra.gov.au

Keep in mind that information given to you by a person or company

promoting or selling a product or therapy may not be reliable, or they may have a financial incentive to recommend a specific treatment. If you have any concerns check with your doctor or pharmacist.

Warning signs

Be on the lookout for the following warning signs when considering any new treatment:

- A cure is offered. There is currently no cure for most forms of arthritis so be wary of products or treatments that promise a cure
- The cure, treatment or therapy is described as secret. Legitimate practitioners share their knowledge so that their peers can review their findings
- Proof for the treatment relies only on testimonials (personal stories). This may be a sign that the treatment has not been scientifically tested
- You are told to give up your current effective treatments or discouraged from getting treatment from your doctor
- The treatment is expensive and not covered by any health fund.

Be wary of products and/or therapies that:

- claim to work for all types of arthritis and/or other disorders
- claim to be free from all side effects
- don't list the ingredients.

If you are still unsure about a treatment for arthritis ask your doctor or pharmacist for advice.

